

## SHORT COMMUNICATIONS

*Acta Cryst.* (1996). **A52**, 955–956

### Vector algebra and crystallography

JEAN SIVARDIÈRE at CEA/Département de Recherche Fondamentale sur la Matière Condensée/MRS, 38054 Grenoble CEDEX 9, France

(Received 15 May 1996; accepted 2 August 1996)

#### Abstract

The volumes of a crystal unit cell and of its reciprocal cell, the relationships between direct and reciprocal interaxial angles and the coordinates of the reciprocal vectors in the direct basis are rederived in a concise way by means of an elementary formula in vector algebra. The product of two rotations is also considered.

The volumes of the direct and reciprocal cells of a crystal and the relationships between direct- and reciprocal-lattice quantities are classically derived using the algebra of determinants and spherical crystallography or advanced vector methods such as Lagrange formulas for products of four vectors. We present here a simple vector formula from which all these quantities are then rederived. Finally, the relationships between direct- and reciprocal-lattice quantities are applied to spherical trigonometry and the product of two rotations.

Consider three linearly independent vectors  $\mathbf{a}$ ,  $\mathbf{b}$ ,  $\mathbf{c}$  and introduce the coordinates  $x, y, z$  of  $\mathbf{c}$  in the  $\mathbf{a}, \mathbf{b}, \mathbf{a} \times \mathbf{b}$  basis:

$$\mathbf{c} = x\mathbf{a} + y\mathbf{b} + z\mathbf{a} \times \mathbf{b}.$$

Taking the scalar product of this relation successively by  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{a} \times \mathbf{b}$ , we find:

$$\mathbf{a} \cdot \mathbf{c} = xa^2 + y\mathbf{a} \cdot \mathbf{b}$$

$$\mathbf{b} \cdot \mathbf{c} = xa \cdot \mathbf{b} + yb^2$$

$$(\mathbf{a}, \mathbf{b}, \mathbf{c}) = z|\mathbf{a} \times \mathbf{b}|^2.$$

Using the Lagrange identity

$$|\mathbf{a} \times \mathbf{b}|^2 = a^2b^2 - (\mathbf{a} \cdot \mathbf{b})^2,$$

we get finally the 'basic' formula

$$|\mathbf{a} \times \mathbf{b}|^2 \mathbf{c} = [b^2(\mathbf{a} \cdot \mathbf{c}) - (\mathbf{a} \cdot \mathbf{b})(\mathbf{b} \cdot \mathbf{c})]\mathbf{a} + [a^2(\mathbf{b} \cdot \mathbf{c}) - (\mathbf{a} \cdot \mathbf{b})(\mathbf{a} \cdot \mathbf{c})]\mathbf{b} + (\mathbf{a}, \mathbf{b}, \mathbf{c})\mathbf{a} \times \mathbf{b}. \quad (1)$$

(a) Suppose that the three vectors  $\mathbf{a}, \mathbf{b}, \mathbf{c}$  define a crystal unit cell: the cell edges are  $a, b, c$  and the interaxial angles are  $\alpha, \beta, \gamma$ . The volume  $V$  of the cell is equal to the triple scalar product  $(\mathbf{a}, \mathbf{b}, \mathbf{c}) = \mathbf{a} \times \mathbf{b} \cdot \mathbf{c} = \mathbf{a} \cdot \mathbf{b} \times \mathbf{c}$ . Taking the scalar product of (1) by  $\mathbf{c}$ , we get the classical expression (Buerger, 1942; Carpenter, 1969; Neustadt & Cagle, 1968; Woolfson, 1970):

$$v^2 = a^2b^2c^2(1 + 2\cos\alpha\cos\beta\cos\gamma - \cos^2\alpha - \cos^2\beta - \cos^2\gamma), \quad (2)$$

which is generally derived using the theory of determinants. An interesting equivalent expression is (Donnay & Donnay, 1959)

$$v^2 = 4a^2b^2c^2[\sin s \sin(s - \alpha) \sin(s - \beta) \sin(s - \gamma)]$$

with  $2s = \alpha + \beta + \gamma$ .

(b) Consider now the reciprocal vectors  $\mathbf{a}^*, \mathbf{b}^*, \mathbf{c}^*$ . We want to determine the reciprocal-cell edges  $a^*, b^*, c^*$ , the interaxial angles  $\alpha^*, \beta^*, \gamma^*$  and the coordinates of the reciprocal vectors in the  $\mathbf{a}, \mathbf{b}, \mathbf{c}$  basis. Since  $v\mathbf{c}^* = \mathbf{a} \times \mathbf{b}$ , we have  $\mathbf{c}^* = ab \sin \gamma / v$ . Equation (1) gives immediately the expression for  $\mathbf{c}^*$  in the direct basis:

$$v^2\mathbf{c}^* = [(\mathbf{a} \cdot \mathbf{b})(\mathbf{b} \cdot \mathbf{c}) - b^2(\mathbf{a} \cdot \mathbf{c})]\mathbf{a} + [(\mathbf{a} \cdot \mathbf{b})(\mathbf{a} \cdot \mathbf{c}) - a^2(\mathbf{b} \cdot \mathbf{c})]\mathbf{b} + |\mathbf{a} \times \mathbf{b}|^2\mathbf{c} \quad (3)$$

or

$$v^2\mathbf{c}^* = ab^2c(\cos\alpha\cos\gamma - \cos\beta)\mathbf{a} + a^2bc(\cos\beta\cos\gamma - \cos\alpha)\mathbf{b} + a^2b^2\sin^2\gamma\mathbf{c}. \quad (3')$$

Comparing this with the general expression

$$\mathbf{c}^* = (\mathbf{a}^* \cdot \mathbf{c}^*)\mathbf{a} + (\mathbf{b}^* \cdot \mathbf{c}^*)\mathbf{b} + (\mathbf{c}^* \cdot \mathbf{c}^*)\mathbf{c} \quad (4)$$

and equating the coefficients of  $\mathbf{b}$  in (3) and (4), we get the angle  $\alpha^*$ :

$$\cos\alpha^* = (\cos\beta\cos\gamma - \cos\alpha) / \sin\beta\sin\gamma \quad (5)$$

Combining (3') and (5) gives

$$v^2\mathbf{c}^* = a^2bc\sin\alpha\sin\gamma\cos\beta^*\mathbf{a} + ab^2c\sin\beta\sin\gamma\cos\alpha^*\mathbf{b} + a^2b^2\sin^2\gamma\mathbf{c}. \quad (6)$$

Calculating  $\sin\alpha^*$  from (5) and (2) gives

$$v = abc\sin\beta\sin\gamma\sin\alpha^* \quad (7)$$

and the sine relation

$$\frac{\sin\alpha}{\sin\alpha^*} = \frac{\sin\beta}{\sin\beta^*} = \frac{\sin\gamma}{\sin\gamma^*}. \quad (8)$$

The volume  $v^*$  of the reciprocal cell is given by an expression similar to (7) and, from (8), one gets  $vv^* = 1$ .

(c) We translate now some of the preceding results into the language of spherical trigonometry.  $\cos\gamma^*$  is given by a relation similar to (5) and the reciprocal relation is

$$\cos\gamma^* = \cos\alpha^*\cos\beta^* - \sin\alpha^*\sin\beta^*\cos\gamma. \quad (9)$$

Using (7) in the form  $v = abc\sin\alpha\sin\beta\sin\gamma^*$  and (8), we may write (6) as

$$abc\sin\alpha^*\sin\beta^*\mathbf{a} \times \mathbf{b} = a^2bc\sin\alpha^*\cos\beta^*\mathbf{a} + ab^2c\sin\beta^*\cos\alpha^*\mathbf{b} + a^2b^2\sin\gamma^*\mathbf{c}. \quad (10)$$

Suppose that the extremities  $A, B$  and  $C$  of the three vectors  $\mathbf{a}, \mathbf{b}, \mathbf{c}$  are on a unit sphere centered at the origin:

$a = b = c = 1$ . The side lengths of the spherical triangle are equal to the interaxial angles  $\alpha, \beta, \gamma$  and its dihedral angles are  $\pi - \alpha^*, \pi - \beta^*, \pi - \gamma^*$ . Using now the classical notations  $a, b, c$  for the side lengths and  $\alpha, \beta, \gamma$  for the dihedral angles, we get from (9) and (10), respectively,

$$\cos \gamma = -\cos \alpha \cos \beta + \sin \alpha \sin \beta \cos c \quad (11)$$

$$\sin \gamma \mathbf{c} = \sin \alpha \cos \beta \mathbf{a} + \cos \alpha \sin \beta \mathbf{b} + \sin \alpha \sin \beta \mathbf{a} \times \mathbf{b}. \quad (12)$$

These formulas were found by Altmann by a more sophisticated procedure (Altmann, 1986). They can be used to derive the product of two rotations around intersecting axes from the Euler–Rodrigues–Hamilton theorem (Altmann, 1986; Sivardière, 1994, 1995). According to this theorem, a rotation of angle  $\theta_1 = 2\alpha$  around  $\mathbf{u}_1 = \mathbf{a}$  followed by a rotation of angle  $\theta_2 = 2\beta$  around  $\mathbf{u}_2 = \mathbf{b}$  is a rotation of angle  $\theta_3 = 2(\pi - \gamma)$  around the unit vector  $\mathbf{u}_3 = \mathbf{c}$ . Introducing the Euler vectors  $\mathbf{R}_i = \sin(\theta_i/2) \mathbf{u}_i$ , we obtain

$$\begin{aligned} \cos(\theta_3/2) &= \cos(\theta_1/2) \cos(\theta_2/2) - \mathbf{R}_1 \cdot \mathbf{R}_2 \\ \mathbf{R}_3 &= \cos(\theta_2/2) \mathbf{R}_1 + \cos(\theta_1/2) \mathbf{R}_2 - \mathbf{R}_1 \times \mathbf{R}_2. \end{aligned}$$

#### References

- Altmann, S. L. (1986). *Rotations, Quaternions and Double Groups*. Oxford: Clarendon Press.
- Buerger, M. J. (1942). *X-ray Crystallography*. New York: John Wiley.
- Carpenter, G. B. (1969). *Principles of Crystal Structure Determination*. New York: Benjamin.
- Donnay, J. D. H. & Donnay, G. (1959). In *International Tables for X-ray Crystallography*, Vol. II. Birmingham: Kynoch Press.
- Neustadt, R. J. & Cagle, F. W. (1968). *Acta Cryst.* **A24**, 247–248.
- Sivardière, J. (1994). *Am. J. Phys.* **62**, 737–743.
- Sivardière, J. (1995). *La Symétrie en Mathématiques, Physique et Chimie*. Grenoble: Presses Universitaires de Grenoble.
- Woolfson, M. M. (1970). *An Introduction to X-ray Crystallography*. Cambridge University Press.